



## MALIGNANT THYROID DISEASE; HYPOCALCAEMIA FOLLOWING THYROID SURGERY

Altaf Ahmed Talpur<sup>1</sup>, Bheesham Kumar<sup>2</sup>, Ahsan Ali Laghari<sup>3</sup>, Abdul Qadeer Shikh<sup>4</sup>, Afzal Junejo<sup>5</sup>, Ali Akber Ghumro<sup>6</sup>

1. FCPS  
Associate Professor  
Surgery unit – IV  
LUMHS, Jamshoro.
2. MBBS,  
Senior Medical Officer  
NMC Specialty Hospital Abu Dhabi,  
UAE.
3. FCPS, MS  
Assistant Professor Surgery  
LUMHS, Jamshoro.
4. MBBS,  
MPH student  
Icahn School of Medicine  
Mount Sinai, Newyork, USA.
5. FCPS (Professor)  
Incharge Surgical Unit – IV  
LUMHS Jamshoro.
6. FCPS (General Surgery)  
Ex-Professor of Surgery  
Peoples Medical College,  
Nawabshah.

### Correspondence Address:

Dr. Altaf Ahmed Talpur  
Flat No: 504, Shaia's Apartments  
Main Jamshoro Road, off Citizen  
Colony  
Hyderabad, Pakistan.  
altafktalpur@yahoo.com

### Article received on:

20/02/2017

### Accepted for publication:

15/09/2017

### Received after proof reading:

29/11/2017

## INTRODUCTION

Tetany and recurrent laryngeal nerve injury are the two most common & ancient complications of thyroid surgery.<sup>1</sup> Initially clinical features of Hypocalcaemia were considered to be due to deficiency of thyroid hormones secondary to thyroid gland surgery. However it was diagnosed as features secondary to Hypocalcaemia due to Parathyroid deficiency when these patients were treated with extracts of parathyroid gland. In their studies they reported 24% of their cases developed Hypocalcaemia due to deficiency of Parathyroid hormone following thyroid surgery.<sup>2</sup>

Immediate clinical manifestations of hypocalcaemia include nerve and muscle

**ABSTRACT... Objectives:** To determine frequency of Transient & permanent hypocalcemia after Thyroid Surgery for Malignant thyroid disease. **Study Design:** Observational study. **Setting:** Public & Private Sector Hospitals of Hyderabad. **Period:** December 2008 to April 2016. **Materials and Methods:** All patients of Thyroid pathology who fulfilled the inclusion criteria were admitted in the ward. They were evaluated preoperatively & surgery was performed. Postoperatively patients were assessed clinically & biochemically for Hypocalcaemia. Patient's data was recorded & analyzed for variables like age, sex, diagnosis of thyroid disease on FNAC, Type of thyroid Malignancy, type of thyroid surgery, Transient & permanent hypocalcemia in relation to type of type of thyroid surgery performed & the hospital stay. **Results:** Total 254 patients were operated for different thyroid pathologies. It includes 91 (35.82%) male & and 163 (64.17%) female patients making ratio of 1: 1.79. Mean age was 37.29 ±7.4 years. Amongst them 33 patients were diagnosed as Malignant Thyroid diseases. Most common type of thyroid malignancy detected was papillary carcinoma in 48.48% patients followed by follicular carcinoma in 42.42% patients. Most common surgical procedures performed includes completion thyroidectomy in 57.57% patients with 01 (3.03%) of them underwent cervical neck dissection followed by total thyroidectomy in 36.36% patients with 02 (6.06%) of them underwent cervical neck dissection. Postoperative hypocalcemia was noted in 45.45% patients with 36.36% patients developed hypocalcemia within 24 hours of operation, 6.06% within 24 to 48 hours & 3.03% patients after 48 hours of surgery. Transient hypocalcemia was noticed in 13/33 (39.39%) cases & permanent hypocalcemia in 02/33 (6.06%) patients. Mean hospital stay was 4.13 ± 0.32 days. **Conclusion:** Hypocalcaemia was noted in 15 (45.45%) patients operated for thyroid Malignancy. Revision surgery & cervical lymph node dissection were noted as risk factors.

**Key words:** Thyroid Malignancy/Surgery/Hypocalcemia.

**Article Citation:** Talpur AA, Kumar B, Laghari AA, Shikh AQ, Junejo A, Ghumro AA. Malignant thyroid disease; hypocalcaemia following thyroid surgery. Professional Med J 2017;24(12):1794-1800. DOI:10.17957/TPMJ/17.3894

symptoms but occasionally psychotic symptoms may be observed. Persistent hypocalcaemia causes disability and a considerable financial loss due to adverse impact on health of patient.<sup>3</sup> Delbridge L described technique of capsular dissection of thyroid gland to reduce the chances of permanent hypocalcemia.<sup>4</sup> Puzziello A et al in their study on 2631 patients reported 0.4% patients operated for thyroid developed permanent hypocalcemia.<sup>5</sup>

Intra operative assay of parathyroid hormone (PTH) predicts risk of hypocalcemia successfully after thyroid surgery as reported by Wang J et al in their study.<sup>6</sup>

Noticeable risk factors of hypocalcemia following thyroid surgery include; large goiters, recurrent goiter, retrosternal goiter extension, hyperthyroidism, total thyroidectomy and experience of the surgeons.<sup>7,8</sup>

The present study was intended to find out the frequency of Hypocalcemia in patients operated for malignant thyroid surgery. It includes risk associated with the extent of surgery & severity of Hypocalcemia.

## MATERIALS & METHOD

The present observational study was conducted at public & private sector hospitals of Hyderabad from December 2008 to April 2016. It includes all patients of either sex above the age of 13 years who underwent different surgical procedures for malignant thyroid diseases. All patients of both sex with age under 13 years, or patients with inoperable thyroid malignancy like stage III or IV thyroid malignancy or patients having clinical or subclinical symptoms & signs of Hypocalcaemia or biochemical measurement show low levels of serum calcium preoperatively or altered serum Albumin levels were excluded. Patients receiving Ca++ supplements, multivitamins and mineral formulas were also excluded. Patient's didn't attend follow-up clinics or consented for study was also excluded.

Pre-operative clinical work up included patient's history, examination of thyroid gland & Neck. Patients were also assessed clinically for symptoms & signs of hypocalcemia. Investigations performed include thyroid hormone profile, serum Ca++ level, serum albumin level, ultrasound of Neck & thyroid gland and FNAC of thyroid swelling or where required thyroid scan was performed. In cases with proven preoperative malignancy of thyroid staging workup was also performed. Indirect laryngoscopy (IDL) was performed preoperatively for vocal cord movement. Diagnosis was made.

Thyroid surgery was planned by surgical team under general anesthesia. Lateral mobilization was performed by capsular dissection technique for parathyroid glands and recurrent laryngeal

nerve. Cervical Lymph nodes dissection was made as necessary depending upon the type of Malignancy & presence or absence of enlarged cervical lymph nodes. Postoperatively patients were assessed clinically for circumoral tingling, numbness of fingers & toes, apprehension, seizures & carpopedal spasm as well as biochemically for hypocalcemia. Postoperatively serum calcium level measurement was performed on all patients on first postoperative day & afterwards if patient developed clinical features of Hypocalcemia on subsequent days. If patient developed clinical features of Hypocalcemia on same postoperative day, serum calcium measurement was performed in that patient on same day. Hypocalcaemia was defined as calcium level <8 mg/dl. If hypocalcemia was confirmed postoperatively, it was treated as per standard protocol. In severe cases 40 MEq of intravenous calcium gluconate diluted in 100 of normal saline was administered. Follow-up visits were advised at 01 week, 01 month for 03 months & then every 03 months for 01 year. Patients were briefed about the diagnosis and procedure to be performed. Patients were informed and permission was granted. They were assured that their participation is voluntary with no harms to them in terms of getting due treatment. They were also given right to withdraw from study without putting any reasons.

Patient's data was recorded on a preformed proforma. Data was analyzed for categorical and continuous variables like age, sex, diagnosis of thyroid disease on FNAC, Type of thyroid Malignancy, type of thyroid surgery, Transient & permanent hypocalcemia in relation to type of type of thyroid surgery performed & the hospital stay. Specific statistical tests were applied & P-value was calculated. P-value less than 0.05 were considered significant. All analysis was done using SPSS 16.

## RESULTS

During this period, total of 254 patients were operated for different thyroid pathologies. It includes 91 (35.82%) male & and 163 (64.17%) female patients making male to female ratio of 1: 1.79. Mean age of this group of patients was

37.29 years with SD of  $\pm 7.4$  years & range of 24 – 67 years.

Amongst 254 operated patients for thyroid pathology, 33 patients were diagnosed as cases of Malignant Thyroid diseases. It includes 13(39.39%) patients who were diagnosed as cases of papillary carcinoma of thyroid on FNAC preoperatively while remaining 20(60.60%) patients were found as cases of Malignant Thyroid diseases on histopathological evaluation of thyroid specimens postoperatively.

Operative procedures performed on 13(39/39%) patients with preoperative diagnosis of Papillary carcinoma of thyroid on FNAC include Total thyroidectomy in 10 (30.30%) patients, total thyroidectomy with cervical lymph node dissection in 02(6.06%) patients while in 01(3.03%) patient thyroid Lobectomy was performed as tumor size was less than 1.5 cm on pre & per operative evaluation. Biopsy of resected specimens confirmed diagnosis of papillary carcinoma of thyroid.

In remaining 20(60.60%) patients, Thyroid Lobectomy was performed as initial procedure as diagnosis was not confirmed on FNAC & specimens were sent for histopathology. It confirmed diagnosis of follicular carcinoma in 14(42.42%) patients, papillary carcinoma in 03(9.09%) patients, Hurthle cell variant of follicular neoplasm in 02(6.06%) patients & medullary carcinoma in 01(3.03%) patient. All these patients underwent completion thyroidectomy except 01(3.03%) patient of Papillary carcinoma in which no further procedure was performed due to small size of tumor i-e less than 1.5 cm. Also 01(3.03%) patient of papillary carcinoma thyroid underwent cervical lymph node dissection along with completion thyroidectomy.

Decreased serum calcium level was noted in 15(45.45%) patients with 11(33.33%) of them had evident clinical features of Hypocalcemia, 03(9.09%) had incipient Hypocalcemia which appeared on application of different clinical signs on 1st Postoperative day & afterwards. While remaining 01(3.03%) patient had no

clinical symptoms or a sign of hypocalcemia but hypocalcemia was recognized on measurement of serum calcium level on 1<sup>st</sup> postoperative day as routine. Amongst the 14(42.42%) patients who developed evident or incipient clinical features of Hypocalcemia, 10(30.30%) developed within 24 hours with 02(6.06%) of them developed this within 06 hours after surgery, 03(9.09%) patients within 24 to 48 hours & 01(3.03%) patients after 48 hours.

All patients with decreased calcium level were put on calcium & vitamin D3 therapy. 11(33.33%) patients with evident clinical features of Hypocalcemia were initially put on intravenous calcium treatment & then switched to oral treatment once clinical features of hypocalcemia settled down. Remaining 04(12.12%) patients of asymptomatic hypocalcemia were kept on oral therapy of calcium. All these patients were also given vitamin D3 therapy.

13(39.39%) patients responded to therapy & on 05<sup>th</sup> postoperative day calcium therapy was stopped. However 02(6.06%) of them developed recurrent hypocalcemia & were put on oral calcium & vitamin D3 therapy until they settled down. In 02(6.06%) patients, clinical features of hypocalcemia did not settled down on calcium & vitamin D3 supplement for 06 weeks consecutively & were declared as cases of permanent hypocalcemia. These 02(6.06%) patients include 01(3.03%) patient each of total thyroidectomy & completion thyroidectomy with cervical lymph node dissection.

Mean duration of hospital stay was  $4.13 \pm 0.32$  days.

Type of Thyroid operation	No: & % of patients
Total Thyroidectomy	10 (30.30 %)
Total Thyroidectomy with Cervical lymph nodes dissection	02 (6.06 %)
Thyroid Lobectomy	0 2 (6.06%)
Completion Thyroidectomy	18 (54.54%)
Completion Thyroidectomy with cervical lymph nodes dissection	01(3.03%)
Total	33

**Table I. Different types of Thyroid operations in Patients with Malignant Thyroid disease (n=33)**

Type of operation	No: & % of patients developed Hypocalcemia			
	Within 24 hours	24- 48 hours	After 48 hours	Total
Total Thyroidectomy(10)	04(12.12%)	01(3.03%)	00	05(15.15%)
Total Thyroidectomy with cervical lymph node dissection (02)	01(3.03%)	00	00	01(3.03%)
Completion Thyroidectomy(18)	06(18.18%)	01(3.03%)	01(3.03%)	08(24.24%)
Completion Thyroidectomy with cervical lymph node dissection(01)	01(3.03%)	00	00	01(3.03%)
Thyroid Lobectomy(02)	00	00	00	00
Total (33)	12(36.36%)	02(6.06%)	01(3.03%)	15(45.45%)

**Table II. Postoperative Hypocalcemia within 24 hours, between 24 to 48 hours & after 48 hours of thyroid surgery According to type of operation**

## DISCUSSION

Present study is one of the recently conducted and reported studies on frequency of hypocalcemia after thyroid surgery in malignant thyroid disease & it was found to be 45.45% from Liaquat University Hospital. Most common type of thyroid malignancy detected was papillary carcinoma in 48.48% patients followed by follicular carcinoma in 42.42% patients. In this study hypocalcemia was noted in 45.45% patients operated for malignant thyroid disorders.

Rosato L et al<sup>9</sup> from Italy mentioned analysis of multicentric study of 14,934 cases of thyroid surgery operated over the period of 05 years. It includes 13% patients with malignant thyroid pathology with 91% belonged to differentiated thyroid carcinoma, 5% to Medullary carcinoma and 4% to undifferentiated carcinoma patients. Various studies from India showed that 76% of thyroid surgeries are conducted for the benign thyroid disorders such as nodular goiters, recurrent nodular goiters, retrosternal goiters, toxic Multinodular goiter and Grave`s disease.<sup>10</sup>

Baldassarre RL et al<sup>11</sup> in their nationwide study of 119,567 thyroidectomy patients from 1998–2008 mentioned postoperative hypocalcemia in 5.5% of all thyroidectomy patients before discharge. They include patients of all thyroid pathologies, not just malignant thyroid. Study by Rosato L et al<sup>9</sup> noticed postoperative transient hypocalcemia in 11.50% patients with malignant thyroid disease with permanent hypocalcemia was seen in 3.25% patients. Study by Page C et al<sup>12</sup> in their study of 351 cases who underwent total thyroidectomy have reported postoperative hypocalcemia in 0.5% to 75% of the patients.<sup>12</sup> Goncalves AJ et

al<sup>13</sup> in their study at Cancer Hospital in Brazil mentioned transient hypocalcemia in 13.1% patients & permanent hypocalcemia in 2.5% patients among 1020 patients who underwent thyroidectomy.

Post-operative hypocalcemia may turn into a serious complication requiring close monitoring and calcium infusions to get rid of the clinical symptoms. Symptoms of post-operative hypocalcemia usually occur in the first 24 to 48 hours.<sup>3,8</sup> Improvements in surgical technique have significantly decreased the complications associated total thyroidectomy. Despite it, post-operative hypocalcemia is still regarded as frequently encountered complication presenting as Tetany in the postoperative period.<sup>7</sup> Literature also mention the experience of surgeon, retrosternal extension of thyroid, type of thyroidectomy, neck dissection and paratracheal lymph node dissection as key factors for postoperative hypocalcemia as mentioned by Mirkin ND et al<sup>14</sup> in their study.

Most common surgical procedures performed in this study includes completion thyroidectomy in 19(57.57%) patients with 01(3.03%) of them also underwent cervical neck dissection followed by total thyroidectomy in 12(36.36%) patients with 02(6.06%) of them underwent cervical neck dissection. Hypocalcemia was noticed in 08/18(44.44%) patients of completion thyroidectomy, in 05/10(50%) patients of total thyroidectomy, 02/03(66.66%) patients who underwent cervical lymph node dissection along with total & completion thyroidectomy indicating extensive surgery poses higher risk of developing hypocalcemia. Of the 02(6.06%)

patients of papillary carcinoma thyroid who underwent Lobectomy, none of them developed hypocalcemia.

In their study by Rosato L et al<sup>9</sup> they mentioned 9599 (64.3%) total thyroidectomies, 3130 (20.9%) total lobectomies, 1448 (9.7%) subtotal thyroidectomies with a monolateral remnant, and 757 (5.1%) subtotal thyroidectomies with bilateral remnants. Among all the patients, 14,057 (94%) were operated on for the first time, and there were 877 (6%) reoperations. They noticed symptomatic hypocalcemia in 10.0% with 8.3% had transient & 1.7% had definitive hypocalcemia which accounts for 63% of all complications. Amongst them transient & permanent Hypocalcemia was seen in 0.4% & 0.07% patients after Lobectomy, in 14% & 2.2% cases after total thyroidectomy and in 5% & 0.6% & 0.8% patients after subtotal thyroidectomies with a monolateral remnant and subtotal thyroidectomies with bilateral remnants respectively. The incidence of postoperative permanent hypocalcemia after malignant thyroid cancer in their study was significantly higher (3.3%).

Iqbal et al<sup>15</sup> from study in Pakistan on patients operated for malignant thyroid diseases reported hypocalcemia in 21.6% patients after total thyroidectomy. Study by Rosato L et al<sup>9</sup> mentioned 11.75% & 2.25% patients developed transient & permanent hypocalcemia after total thyroidectomy, 0.5% & 0 after Lobectomy, 4.6% & 0.4% after subtotal thyroidectomy with monolateral residue & 4.1% & 0.9% after subtotal thyroidectomy with bilateral residue. Study by Tredici P et al<sup>16</sup> from Italy mentioned postoperative hypocalcemia in 9.0% patients with total thyroidectomy, in 1.9% following unilateral thyroid Lobectomy, 14.4% following thyroidectomy plus unilateral neck dissection, 23.4% following thyroidectomy plus bilateral neck dissection, 9.6% following complete substernal thyroidectomy, 3.4% following partial substernal thyroidectomy, 6.5% following substernal thyroidectomy, and 3.4% following isthmectomy or otherwise unspecified partial thyroidectomy. Goncalves AJ et al<sup>13</sup> in their study at Sao Paulo mentioned unilateral subtotal thyroidectomy

associated with contralateral hemithyroidectomy was the most prevalent surgical resection causing hypocalcemia. It occurred in 66.6% of the cases followed by total thyroidectomy in 53.4% and by bilateral subtotal thyroidectomy in 35.3% cases. In their study the incidence of hypocalcemia in patients with malignant disease was higher (60.7 percent) than those with benign disease (46.4 percent). Jisheng Hu et al<sup>17</sup> in their study on 5559 patients mentioned correlative incidence of transient & permanent hypocalcemia in patients who were operated for first time of thyroid surgery & in patients who underwent revision surgery. They found temporary hypocalcemia in 3.53% patients & permanent hypocalcemia in 0.02% patients in first time surgery on thyroid while 13.76% & 1.59% patients developed transient & permanent hypocalcemia respectively in revised surgery group. In first time surgery group, transient & permanent hypocalcemia was noticed in 3.28% & 0.02% patients after total thyroidectomy, 2.18% & 0 in patients of hemithyroidectomy, 8.75% & 0 in patients of total thyroidectomy with central neck dissection group & 9.26% & 0.02% patients of total thyroidectomy with mono neck dissection. In the revised surgery group transient hypocalcemia & permanent hypocalcemia was noticed in 1.38% & 2.17% patients who underwent completion thyroidectomy after subtotal thyroidectomy & in 1.96% & 0 patients who underwent contralateral thyroidectomy after hemithyroidectomy at first surgery. Jong-Lyel R et al<sup>18</sup> in their study mentioned morbidity and hypocalcemia was higher in the node dissection group than the no node dissection group & serum PTH levels significantly decreased immediately postoperatively in the node dissection group and remained low for several weeks thereafter.

In this study 36.36% patients developed hypocalcemia within 24 hours of operation on thyroid, 6.06% within 24 to 48 hours of surgery & 3.03% patients after 48 hours of surgery. Mirkin ND et al<sup>14</sup> in their study reported Hypocalcemia in 54% patients within the first 24 hours postoperatively with 40% of them had symptomatic hypocalcemia. Study by Nair et al<sup>3</sup> has reported a frequency of Hypocalcemia in 23% subjects. Nair et al<sup>3</sup> further reported onset of hypocalcaemia within 6 hours

in 11.05% patients and was delayed up to 3<sup>rd</sup> post operative day in 6.84%. Tredici P et al<sup>16</sup> reported that 50(50%) patients of their patients developed postoperative hypocalcemia with 24(48%) of them had symptomatic hypocalcemia & 26(52%) had asymptomatic hypocalcemia. Amongst them 12.5% had on the first day after surgery, 62.5% on the second day, 17% on the third day and 4% each on the fourth and fifth days.

In this study transient hypocalcemia was noticed in 13/33 (39.39%) cases & permanent hypocalcemia in 02/33(6.06%) patients. These 02(6.06%) patients include one patient of total thyroidectomy & other patient of completion thyroidectomy with cervical lymph node dissection. Iqbal et al<sup>15</sup> reported transient hypocalcemia in all cases.

Risk of getting permanent hypocalcemia varies with the type of surgery, expertise of operating surgeon. Lower chances of developing hypocalcemia have been noticed in highly specialized centers of thyroid pathology.

Analysis of Parathyroid hormone level 1 to 6 hours after thyroidectomy has significant accuracy in determining the chances of developing symptoms of hypocalcemia. Its routine measurement may help in earlier discharge of these patients and also early identification of developing hypocalcemia that may be instituted early treatment of post thyroidectomy hypocalcemia.<sup>19</sup>

Jong-Lyel R et al<sup>20</sup> in their study of 197 patients operated for differentiated papillary thyroid carcinoma with 49 of them underwent total thyroidectomy alone and 148 underwent total thyroidectomy plus central neck dissection. Patients with neck dissection were randomized into 03 groups with Group A received oral calcium plus vitamin D, group B received only calcium, Group C received no supplemental therapy. Amongst them Group C patients developed significantly higher incidence of hypocalcemia which can be prevented by routine postoperative supplementation with oral calcium and vitamin D.

In this study mean duration of hospital stay was  $4.13 \pm 0.32$  days while Mirkine ND et al<sup>14</sup> mentioned

this as  $3.33 \pm 3.09$  days for hypocalcemic patients and  $1.85 \pm 3.55$  days for normocalcemic patients. Tredici P et al<sup>16</sup> found hospital stay of 6.19 days  $\pm$  0.24 days.

## CONCLUSION

Relatively low number of Malignant thyroid disease (12.99%) was noticed in this study with incidence of hypocalcemia was 45.45% in all operations performed for malignant thyroid disorder. Repeat surgery on thyroid with more extensive dissection predisposes the patient to develop hypocalcemia.

## Limitations of the study

Non measurement of vitamin D level and parathyroid hormone level due to cost and funding issues.


Copyright© 15 Sep, 2017.

## REFERENCES

- Fortuny VJ, Guigard S, Karenovics W, Triponez R. **Surgery of the thyroid: Recent developments and perspective.** Swiss Medical Weekly 2015; 1451: 141 - 44.
- Michie W, Stowers JM, Frazer SC, Gunn A. **Thyroidectomy and the Parathyroids.** Br J Surg 1965; 52: 503-14.
- Nair CG, Babu MJ, Menon R, Jacob P. **Hypocalcaemia following total thyroidectomy: An analysis of 806 patients.** Indian J Endocr Metab 2013; 17:298-303.
- Delbridge LW. **Total Thyroidectomy: The evolution of surgical technique.** ANZ J Surg 2003; 73:761- 8.
- Puzziello A, Rosato L, Innaro N, Orlando G, Avenia N, Perigli G, Calo PG, et al. **Hypocalcemia following thyroid surgery: incidence and risk factors. A longitudinal multicenter study comprising 2,631 patients.** Endocrine 2014; 47:537-542.
- Wang J, Gu J, Han Q, Wang W, Shang J. **Value of intra operative parathyroid hormone monitoring in papillary thyroid cancer surgery: Can it be used to guide the choice of operation methods?** Int J Clin Exp Med 2015; 8(5):7778-7785.
- Bliss RD, Gauger PG, Delbridge LW. **Surgeon's approach to thyroid gland- Anatomy and surgical technique.** World J Surg 2011; 24:891- 7.
- Salinger EM and Moore JT. **Perioperative indicators of hypocalcemia in total thyroidectomy: the role of**

- vitamin D and parathyroid hormone.** Am J Surg 2013; 206: 876 - 881.
9. Rosato L, Avenia N, Bernante P, Palma MD, Gulino G, Nasi PG, Pelizzo MR, Pezzullo L. **Complications of Thyroid Surgery: Analysis of a Multicentric Study on 14,934 Patients Operated on in Italy over 5 Years.** World J Surg 2004; 28(3): 271 – 276.
  10. Menon VU, Sundaram KR, Unnikrishnan AG, Jayakumar RV, Nair V, Kumar H. **High prevalence of undetected thyroid disorders in an iodine sufficient adult in south Indian population.** J Indian Med Assoc 2009; 107:72- 7.
  11. Baldassarre RL, Chang DC, Brumund KT, Bouvet M. **Predictors of Hypocalcemia after Thyroidectomy: Results from the Nationwide Inpatient Sample.** ISRN Surgery 2012.1-7. Article ID 838614. <http://dx.doi.org/10.5402/2012/838614>.
  12. Page C, Strunski V. **Parathyroid risk in total thyroidectomy for bilateral, benign, multinodular goiter: Report of 351 surgical cases.** J Laryngol Otol 2007; 121: 237-41.
  13. Gonçalves AJ, Martins L, DeSouza TRB, Alves PJC, Schiola A, Rios OAB, Dos Santos AR. **Clinical and laboratorial correlation of postoperative hypocalcemia after extensive thyroidectomy.** Sao Paulo Med. J 1997; 115(1): 234 - 239. <http://dx.doi.org/10.1590/S1516-31801997000100009>.
  14. Mirkine ND, Hooghe L, Geertruyden JV, Maertelaer VD. **Hypocalcemia after Thyroidectomy. Predictors of hypocalcemia occurring after a total/near total thyroidectomy.** Arch Surg 1992; 127(7): 854-858.
  15. Iqbal M, Subhan A, Baig MS, Shah MS. **Frequency of hypocalcemia in total thyroidectomy.** J Surg Pak 2010; 15 (2):87-91.
  16. Tredici EP, Grosso , Gibelli MA, Massaro C, Arrigoni N, Tradati N. **Identification of patients at high risk for hypocalcemia after total thyroidectomy.** Acta Otorhinolaryngol Ital. 2010; 30: 144-148. PMID: PMC3185817.
  17. Jisheng H, Zhao N, Kong R, Wang D, Sun B, Wu L. **Total thyroidectomy as primary surgical management for thyroid disease: surgical therapy experience from 5559 thyroidectomies in a less-developed region.** World J Surg Onco 2016; 14(20): 1-7. DOI 10.1186/s12957-016-0772-1.
  18. Jong-Lyel R, Jae-Yong P, Chan II PMD. **Total Thyroidectomy plus Neck Dissection in Differentiated Papillary Thyroid Carcinoma Patients: Pattern of Nodal Metastasis, Morbidity, Recurrence, and Postoperative Levels of Serum Parathyroid Hormone.** Annals of Surgery: April 2007; 245(4): 604-610. doi: 10.1097/01.sla.0000250451.59685.67.
  19. Noordzij JP, Lee SL, Bernet VJ, Payne RJ, Cohen SM, McLeod IK, Hier MP, Black MJ, Kerr PD, Richards ML, Lo CY, Raffaelli M, Bellantone R, Lombardi CP, Cohen JI, Dietrich MS. **Early Prediction of Hypocalcemia after Thyroidectomy using Parathyroid Hormone: An Analysis of Pooled Individual Patient Data from Nine Observational Studies** J Am coll Surg 2007; 06: .298-302. DOI: <http://dx.doi.org/10.1016/j>.
  20. Jong-Lyel R, Jae-Yong P, Chan-II P. **Prevention of postoperative hypocalcemia with routine oral calcium and vitamin D supplements in patients with differentiated papillary thyroid carcinoma undergoing total thyroidectomy plus central neck dissection.** Cancer: 2009; 115(2): 251–258.

### AUTHORSHIP AND CONTRIBUTION DECLARATION

Sr. #	Author-s Full Name	Contribution to the paper	Author=s Signature
1	Altaf Ahmed Talpur	Design of the work & acquisition of data	
2	Bheesham Kumar	Analysis & interpretation of data	
3	Ahsan Ali Laghari	Drafting the article	
4	Abdul Qadeer Shikh	Recising critically for important intellectual content	
5	Afzal Junejo	Final approval of the version	
6	Ali Akber Ghumro	Revised & agreed to all aspects of works	