



# ATRIAL FIBRILLATION; TO DETERMINE THE FREQUENCY AND ITS PREDISPOSING FACTORS IN PATIENTS AFTER CORONARY ARTERY BYPASS GRAFTING.

Rehan Riaz<sup>1</sup>, Nabeel Ahmad<sup>2</sup>, Shahid Abbas<sup>3</sup>

1. FCPS (Card)  
Assistant Professor of Cardiology,  
Faisalabad Institute of Cardiology,  
Faisalabad.
2. M.B;B.S.  
Post Graduate Registrar Cardiac  
Surgery,  
Faisalabad Institute of Cardiology  
Faisalabad,
3. FCPS (Card)  
Assistant Professor Cardiology  
Faisalabad Institute of Cardiology,  
Faisalabad, Pakistan.

**Corresponding Address:**

Dr Rehan Riaz  
Assistant Professor of Cardiology,  
Faisalabad Institute of Cardiology  
Faisalabad.  
rehanriazdab@gmail.com

**Article received on:**

15/08/2017

**Accepted for publication:**

20/10/2017

**Received after proof reading:**

29/11/2017

**ABSTRACT... Introduction:** Atrial Fibrillation (AF) is one of the most common arrhythmia after coronary artery bypass grafting (CABG). Many risk factors have been identified for the development of postoperative AF with varied level of evidence. AF is associated with increased risk of morbidity and mortality causing prolong hospital stay and utilizing more resources. Our study was aimed to identify high risk population for developing post-operative AF and to draw recommendations for its prevention. **Objectives:** Our objective was to determine the frequency of Atrial Fibrillation (AF) and its predisposing factors in patients after Coronary Artery Bypass Grafting (CABG). **Setting:** The study was conducted at Cardiac Surgery Department / Faisalabad Institute of Cardiology, Faisalabad. **Duration of Study:** 15-12-2014 to 14-07-2015. **Study Design:** Cross sectional study. **Results:** The result showed that out of 130 patients undergoing CABG surgery, 104 (80.3%) male and 26 (19.7%) females, 9(7%) patients suffered AF in post-operative phase. A significant association was found between postoperative AF and advanced age ( $p = 0.011$ ), obesity ( $p = 0.028$ ), low EF ( $p = 0.000$ ), and post-operative use of  $\beta$ -blocker ( $p = 0.007$ ). Significant difference was found between postoperative AF and days stayed in hospital ( $p = 0.000$ ) and magnesium level at first day ( $p = 0.038$ ). On the other hand there was no significant relationship found between AF and prolongs cross clamp and bypass time. **Conclusion:** Advance age, obesity, congestive heart failure, low ejection fraction and withdrawal from beta blockers in post-operative period have been identified as a significant risk factors for the development of post-operative atrial fibrillation.

**Key words:** Atrial Fibrillation, Coronary Artery Bypass Grafting, Postoperative Arrhythmia.

**Article Citation:** Riaz R, Ahmad N, Abbas S. Atrial fibrillation; To determine the frequency and its predisposing factors in patients after coronary artery bypass grafting. Professional Med J 2017;24(12):1852-1859.  
DOI:10.17957/TPMJ/17.4245

## INTRODUCTION

Atrial fibrillation is frequent arrhythmic complications of coronary artery bypass grafting (CABG) as its incidence varies from 20 to 40 % primarily within 2 to 3 days.<sup>1,2</sup> The incidence of AF has been estimated to be 26.7% in a meta-analysis of 24 trails.<sup>3</sup> Over the past decades the incidence of post CABG AF is continuously increasing.<sup>3</sup>

For the development of postoperative AF with varied level of evidence many risk factors have been identified. The strong association of prior AF, advanced age, male gender, obesity, postoperative beta blocker withdrawal have been established, while other factors such as congestive heart failure (CHF), chronic lung disease, systemic emboli, decreased (LV/EF),

increased left atrial (LA) size, hypomagnesaemia, myocardial infarction peri and postoperative, prolong cross clamp and bypass time have shown less likely association with this arrhythmia.<sup>4</sup> These frequent post-operative cardiac arrhythmias linked and consequently prolong the hospital stay. The episodes of AF are usually symptomatic as patients experience palpitation, nausea and malaise.<sup>5</sup>

Identification and thus reduction of predisposing factors may result in better treatment strategies for the common cardiac arrhythmia. Furthermore, few local studies are available which can identify the frequency and the predisposing factors of post-operative AF in our patient population.<sup>6</sup>

The current study was aimed to evaluate the

frequency of Atrial Fibrillation in post CABG and to determine the inclining factors associated with AF after CABG in population understudy.<sup>7</sup> The resulting commendations may aid in reduction of morbidity and mortality as it will enable us to predict the patient may at the risk of developing AF and hence, save them from having consequential arrhythmia. This will lead in minimize the duration of hospital stay and cost, both to the patient and the hospital.<sup>8</sup>

## PATIENTS AND METHODS

### OBJECTIVE

Our objective was to determine the frequency of Atrial Fibrillation (AF) and its predisposing factors in patients after Coronary Artery Bypass Grafting (CABG).

### OPERATIONAL DEFINITIONS

#### Atrial Fibrillation

On the electrocardiogram (ECG) the irregularity of P – waves that vary in timing would be considered as presence of Atrial Fibrillation (AF) in first 3-5 post-operative days of ICU stay.

#### Predisposing Factors

##### Advanced age

Patient with age  $\geq$  50 years was taken as advanced age patients.

##### Obesity

Patient with BMI  $\geq$  30 was taken as obese patients.

##### Decreased LV/EF

Patient with EF  $\leq$  40 % was taken as patients with decreased LV/EF.

##### Hypomagnesaemia

Patients having serum level of magnesium  $\leq$  1.6mmol/L was taken as Hypomagnesaemia.

##### Post-operative withdrawal from $\beta$ - blockers

Patient who were taking beta blockers for more than 6 weeks was selected

### Perioperative Myocardial infarction (MI)

Patients with chest pain and ST elevation more than 2mm in any lead on ECG or with raised cardiac enzymes was taken as patients of perioperative MI.

## MATERIAL AND METHODS

### Setting

The study was conducted in Cardiac Surgery Department / Faisalabad Institute of Cardiology, Faisalabad.

### Duration of Study

15-12-2014 to 14-07-2015

### Study Design

Cross sectional study

### Sampling Technique

Non-probability consecutive sampling.

### Sample Size

130 consecutive Patients  
(The sample size has been calculated using WHO Sample size calculator)  
Absolute precision required 6%, P=13.7%,  
Confidence level 95%

### Sample Selection

### Inclusion Criteria

All Patients undergo electively CABG on both genders in age group 30-80 years.

### Exclusion Criteria

- Patients undergoing for Re – do CABG
- Patients having IHD with concurrent valvular heart disease
- Patient having prior documented history of AF

### Methodology

Informed consent was taken from all the patients fulfilling the inclusion criteria. Patients' clinical data (age, sex) and information regarding factors, which can predispose the patient to AF, these have include (advanced age, male gender, obesity, CHF, chronic lung disease, systemic emboli, decreased LV/EF, hypo-magnesaemia,

peri and post-operative MI, prolong cross clamp and bypass-time) were recorded in the pre-designed Proforma.

Post operatively Patients were monitored in the Cardiac Surgery ICU / Ward information about the presence of factor which predisposes AF was recorded. Occurrence of AF was noted and patients were divided into two groups in the presence or absence of AF.

### Data Collection Tools

Quantitative and qualitative variables were entered into the predesigned proforma which is attached as appendix

### STATISTICAL ANALYSIS

The data was entered and analyzed using SPSS (Statistical Package for social Sciences) 18.0. Mean +SD was calculated for quantitative variables like age. Frequencies and percentage was calculated for qualitative parameters like gender, AF, Obesity, decreased EF, withdrawal from beta blocker, prolong pump or cross clamp time and hypomagnesaemia. Effect modifiers like age and gender was controlled by stratification and post stratification chi-square test was applied. A P – Value of  $\leq 0.05$  was considered as statistically significant.

### RESULTS

The result shows that out of 130 patients undergoing CABG surgery, 104 (80%) male and 26 (20%) were females, 9(7%) patients suffered AF in post-operative phase. The mean and standard deviation of age was observed as  $55.81 \pm 9.00$  years. A significant association was found between postoperative AF and advanced age ( $p = 0.011$ ), obesity ( $p = 0.028$ ), low EF ( $p = 0.000$ ), and post-operative use of  $\beta$ -blocker ( $p = 0.007$ ). Significant difference was found between postoperative AF and days stayed in hospital ( $p = 0.000$ ) and magnesium level at first day ( $p = 0.038$ ). On the other hand there was no significant relationship found between AF and prolonged cross clamp and bypass time. No patient suffered from perioperative MI so no data regarding its relationship could be established with post-operative AF.

A total of 7 male patients and 2 female patients suffered from AF post operatively and there was no significant association found between post CABG AF and gender ( $p=0.475$ ).

All the patients were divided into two groups regarding age. There were 102 (78.4%) patients having age of 50 years or more and 28 (21.6%) patients having age less than 50 years. There was significant association between post operative AF and advanced age ( $p=0.04$ ) as 7 patients (88 %) were in older group ( $>50$  years), while only 2 patients (22%) were in younger group ( $<50$  years).

Statistically significant relationship has been observed between obesity and development of AF post operatively ( $p=0.022$ ) as out of 13 obese patients (BMI  $>30$ ) 4 (11%0 developed AF.

There was no significant difference found between post-operative AF and pre-operative hypomagnesaemia. But there was significant difference found between postoperative AF and magnesium level at first day ( $p=0.017$ ).

A total of 17 patients (13%) had ejection fraction (EF)  $<40\%$  whereas 113 (87%) patients with EF  $>40\%$ . There was significant association observed between post-operative AF and low EF ( $p=0.000$ ), AF occurred in 6 (35.2 %) patients with low EF.

A total of 30 patients (24%) were on Beta blocker pre-operatively and 100 (76%) patients were not taking beta blocker. There was significant association between post CABG AF and use of  $\beta$ -blocker withdrawal ( $p=0.03$ ). AF has occurred in 4 patients out 30 patients using  $\beta$ -blocker pre operatively.

No patient suffered from myocardial infarction peri and post operatively in each condition. So no association can be determined.

Cardiopulmonary bypass (CPB) time ( $>120$  min) was documented in 12 (10.1%) patients, while CPB time less than 120minutes was observed in 118 (90%) patients. This study failed to observe

any statistically significant association between post-operative AF and prolong CPB time ( $p=0.057$ ).

35 (25%) patients required more than 90 minutes of aortic cross clamp time, while the majority of the (75%) 95 patients required less than 90 minutes to complete their grafting. No significant association

was observed between post-operative AF and prolong ACC time ( $p=0.376$ ).

The mean and standard deviation of days stayed in ICU  $5.22 \pm 2.6$ . It is observed that in patients who developed AF the ICU stay as well the hospital stay was prolonged.

Variable	Minimum	Maximum	Mean	Standard Deviation
Age	30	80	55.1	9.002
Height	122	190	164.66	13.13
Weight	35	118	73.6	9.35
BMI	14.3	47.9	27.1	4.56
Magnesium level	1.3	3.4	2.012	0.29

**Table-I. Mean and std. deviation of quantitative variables**

Gender		Frequency		Percentage
Male		104		80.3%
Female		26		19.7%
Total		130		100%
Gender* AF during ICU stay				
		AF during ICU stay		Total
		Yes	No	
Gender	Male	7	97	104
	Female	2	24	26
Total		9	121	130
P value 0.475* Chi Square ( $>0.05$ )** Not Significant				

**Table-II. Distribution of patient by gender (n=130)**

Age in years		Frequency		Percentage
Age 50 years and above		33		25%
Age less than 50 years (30-49)		97		75%
Total		130		100%
Age * AF during ICU stay				
		AF during ICU stay		Total
		Yes	No	
Age	>50 years	7	26	33
	<50 years	2	95	97
Total		9	121	130
P value 0.04* Chi Square ( $>0.05$ )** Significant				

**Table-III. Distribution of patients by age**

Obesity		Frequency		Percentage
BMI more or equal 30		34		26%
BMI less than 30		96		74%
Total		130		100%
Obesity * AF during ICU stay				
		AF during ICU stay		Total
		Yes	No	
Age	BMI >30	4	30	34
	BMI <30	5	91	96
Total		9	121	130
P value 0.022* Chi Square (>0.05)** Significant				

**Table-IV. Distribution of patient by obesity**

**DISCUSSION**

The present research has been conducted at Faisalabad Institute of Cardiology Faisalabad which is one of the largest hospitals only dedicated to deal with patients having heart diseases. AF has been considered as most common arrhythmia to occur after CABG and is associated with increased morbidity and mortality and longer, more expensive hospital stays.<sup>9,10</sup> The incidence of post-operative AF (POAF) varies between 20-40% as reported in different studies. The incidence of POAF has increased continuously over the past decades and it's believed to be either due to aging of the population undergoing heart surgery or deployment of more sensitive methods of surveillance of AF, such as telemetric monitoring.<sup>11,12</sup>

According to MEDLINE database reports published between 2010 and June 2014 by using the keyword atrial fibrillation, coronary artery bypass grafting. The author found 10 to 65 % of patients developing AF post operatively depending on patient's profile, type of surgery, method of arrhythmia surveillance and definition of arrhythmia.<sup>3</sup> Patient undergoing CABG alone have lower incidence of post-operative AF than patients undergoing valve surgery or combined CABG–valve operations. The Meta analysis of 24 trails included in this study estimated the incidence of AF at 26.7% in patients undergoing CABG alone.<sup>4</sup> The present study has also revealed less incidence of AF in our population as 9 out of 130 patients (7 %) developed AF post operatively.

Review of literature has consistently shown the advance age as predisposing factor for the development of AF after CABG. A study suggested that older age is the most effective predisposing reason for AF because ageing may lead to dilatation and an enhancement in fibrosis.<sup>16</sup> It shown significant effect of advanced age for post-operative AF with p <0.001. A similar trend also reported, in which he found the significant association between advance age and post-operative atrial fibrillation (OR for 10-year increase, 1.75; 95% CI, 1.59-1.93). It is also reported that the overall frequency of AF rising 1.8% with age. The incidence of AF in patients <70 years is estimated to be 0.4%, whereas for patients >70 years, the incidence is 2% to 4%.<sup>17</sup> It was searched that based on the patient's age, the frequency of AF deviated from 13% for patients younger than 50 years of age to 52% for those 80 years of age or older, with an OR of 1.60 for each decade of additional life. Some said that older age may cause structural changes in the heart such as an increase in chamber size and fibrosis. They reported that along with these changes other age related comorbidities seems to be responsible for increased incidence of AF with increasing age.<sup>18,19</sup> Consistent with other reports in which older age constantly has been discovered to be a self-determining predictor for post CABG AF, our research also revealed that advance age (more than 50 years) has significant association for the occurrence of post operative AF (95% CI 1.02 to 1.08, 0.04).

The patients having BMI >30kg/m2 undergoing

CABG have consistent reports regarding development of POAF. Studies showed that mild obesity and moderate severe obesity were independent risk factors for POAF in a retrospective study in 4583 patients over 50 years of age, undergoing CABG. Moreover, some found BMI to be an independent predictor for new-onset AF in the general population.<sup>21</sup> Studies found that obesity, age and male gender to be independent predictors of post-operative AF.<sup>13,14,15</sup> Similarly in our study, we have observed the preoperative obesity (body mass index >30 kg/m<sup>2</sup>) in which there were 24.7% patients having obesity. By using chi-square analysis we came to know that obesity have a significant association for post-operative AF having p-value = 0.022, as body mass index enhances the risk of post-operative AF. So obesity is considered as one of the risk factors for the post-operative AF after CABG.

Prior studies have determined that Low left ventricular ejection fraction (LVEF) is an important risk factors which may predispose patients undergoing CABG to develop POAF. Research work reported in their study that low ejection fraction is a significant risk factor, predisposing POAF. Similarly some other studies have also shown the significance of EF <40% regarding POAF. They found that patients with an LVEF <30% predisposes post-operative AF and had slightly increased mortality.<sup>20</sup>

Another risk factor that has been evaluated regarding POAF was beta blocker (BB) withdrawal after surgery. In a meta analysis of thirty one trails evaluated has shown that beta blocker withdrawal predisposing POAF (OR 0.30, 95% CI 0.22 to 0.4). ACC/ AHA quoted that withdrawal of beta-blockers double the risk of AF after CABG. Thus, early re-initiation of beta blockers is critical for avoidance of this complication.<sup>22,23</sup> Similarly we also observed that withdrawn of beta blocker post operatively predisposes the risk of POAF. (OR 1.112, 95% CI 1.16 to 1.24, p=0.03).

It has been proposed that prolong CPB and cross clamp time might impact post CABG AF incidence, but didn't achieve accepted levels for statistical significance (p=0.57) and (p=0.376).

Existing literature also shows that the administration of magnesium level reduces the post-operative risk of AF. Studies have noticed that the management of magnesium level reduces the frequency of AF in patients undergoing CABG.<sup>24</sup> In the present research we found significant difference between the magnesium level at day 1 post-operative AF (p=0.017) which shows that there is significant effect of low magnesium level on post-operative AF. We also observed that there was significant difference found between postoperative AF and days stayed in hospital (p = 0.000) and magnesium level at first day (p = 0.038).

### CONCLUSION


Atrial Fibrillation is common complication following coronary artery bypass grafting surgery. Which not only increases morbidity and mortality but also increase financial burden on account of longer ICU and hospital stay. Advance age, obesity, congestive heart failure, low ejection fraction and dilated LA have been identified as a significant risk factors for the development of post-operative atrial fibrillation. So different modalities such as administration of beta blockers, amiodarone, magnesium, can be utilized to prevent POAF in the groups of high risk patients undergoing CBAG. This will reduce not only morbidity and mortality in these high risk patients but also help to reduce cost of surgery which is of great importance especially in our part of the world.

Copyright© 20 Oct, 2017.

### REFERENCES

1. Filardo G1, Hamilton C, Hebelers RF Jr, Hamman B, Grayburn P. **New-onset postoperative atrial fibrillation after isolated coronary artery bypass graft surgery and long-term survival.** CircCardiovascQual Outcomes, 2009; 2:164-9.
2. Rostagno C, La Meir M, Gelsomino S, Ghilli L, Rossi A, Carone E, et al. **Atrial fibrillation after cardiac surgery: incidence, risk factors, and economic burden.** J CardiothoracVascAnesth 2010; 24:952-8.
3. Shen J, Lall S, Zheng V, Buckley P, Damino RJ. **The persistent problem of new onset postoperative atrial fibrillation: A single- institution experience over two decades.** J ThoracCardiovasc Surg. 2011; 141;559-70.

4. Lapar DJ, Spir AM, Corsby IK, Fonner E, Brown M. **Postoperative atrial fibrillation significantly increases mortality, hospital readmission and hospital cost.** *Ann Thoracic Surg* 2014; 98:527-33.
5. Helgadottir S, Sigurdsson MI, Ingvarsdottir IL, Arnar DO, Gudbjartsson T. **Atrial fibrillation following cardiac surgery: risk analysis and long-term survival.** *J Cardiothorac Surg.* 2012; 7:87.
6. Mostafa A, Mohamed A. **El Hadeed, Shenoy M, Tushartuliani.** *Avicenna J Med.* 2012;2:65-70.
7. Tadic M, Ivanovic B, Zivkovic N, **Predictors of atrial fibrillation following coronary artery bypass surgery.** *Med SciMonit,* 2011; 17:R48-55.
8. Bramer S, van Straten AH, Soliman Hamad MA, Berreklouw E, Martens EJ, Maessen JG. **The impact of new-onset postoperative atrial fibrillation on mortality after coronary artery bypasses grafting.** *Ann Thorac Surg* 2010; 90:443-9.
9. Aranki SF, Shaw DP, Adams DH, Rizzo RJ, Couper GS, VanderVliet M, Collins JJ, Cohn LH, Burstin HR. **Predictors of atrial fibrillation after coronary artery surgery.** *Circulation.* 1996; 94:390-7.
10. Creswell LL, Schuessler RB, Rosenbloom M, Cox JL. **Hazards of postoperative atrial arrhythmias.** *Ann Thorac Surg.* 1993; 56:539-49.
11. Kalman JM, Munawar M, Howes LG, Louis WJ, Buxton BF, Gutteridge G, Tonkin AM. **Atrial fibrillation after coronary artery bypass grafting is associated with sympathetic activation.** *Ann Thorac Surg.* 1995; 60:1709-15.
12. Echahidi N, Mohty D, Pibarot P, Despre's JP, O'Hara G, Champagne J, Philippon F, Daleau P, Voisine P, Mathieu P. **Obesity and metabolic syndrome are independent risk factors for atrial fibrillation after coronary artery bypass graft surgery.** *Circulation.* 2007; 116(11 Suppl): I-213: I-219.
13. Wang TJ, Parise H, Levy D, D'Agostino RB, Sr, Wolf PA, Vasan RS, Benjamin EJ. **Obesity and the risk of new-onset atrial fibrillation.** *JAMA.* 2004; 292(20):2471-7.
14. Pritchett AM, Jacobsen SJ, Mahoney DW, Rodeheffer RJ, Bailey KR, Redfield MM. **Left atrial volume as an index of left atrial size: a population-based study.** *J Am CollCardiol.* 2003; 41(6):1036-43. [PubMed].
15. Zacharias A, Schwann TA, Riordan CJ, Durham SJ, Shah AS, Habib RH. **Obesity and risk of new-onset atrial fibrillation after cardiac surgery.** *Circulation.* 2005; 112(21):3247-55.
16. Omer S, Cornwell L, Bakshi A, et al. **Incidence, Predictors, and Impact of Postoperative Atrial Fibrillation after Coronary Artery Bypass Grafting in Military Veterans.** *Tex Heart Inst J.* 2016; 43(5): 397-403.
17. Mathew JP, Fontes ML, Tudor IC, Ramsay J, Duke P, Mazer CD et al. **A multicenter risk index for atrial fibrillation after cardiac surgery.** *JAMA.* 2004; 291(14):1720-9.
18. Allessie MA, Boyden PA, Camm AJ, Kleber AG, Lab MJ, Legato MJ et al. **Pathophysiology and prevention of atrial fibrillation.** *Circulation.* 2001; 103(5):769-77.
19. Spach MS, Dolber PC, Heidlage JF. **Influence of the passive anisotropic properties on directional differences in propagation following modification of the sodium conductance in human atrial muscle. A model of reentry based on anisotropic discontinuous propagation.** *Circ Res.* 1988; 62(4):811-32.
20. Borde D1, Gandhe U, Hargave N, Pandey K, Mathew M, Joshi S. **Prediction of postoperative atrial fibrillation after coronary artery bypass grafting surgery: is CHA 2 DS 2 -VASc score useful?** *Ann Card Anaesth.* 2014; 17(3):182-7.
21. Bramer S, van Straten AH, Soliman Hamad MA, et al. 2011. **Body mass index predicts new-onset atrial fibrillation after cardiac surgery.** *Eur J CardiothoracSurg;* 40(5):1185-90.
22. Skiba, M., Pick, A., Chaudhuri, K. et al, **Prophylaxis against atrial fibrillation after cardiac surgery: beneficial effect of perioperative metoprolol.** *Heart Lung Circ.* 2013; 22:627-633.
23. Khan, M.F., Wendel, C.S., Movahed, M.R. **Prevention of post-coronary artery bypass grafting (CABG) atrial fibrillation: efficacy of prophylactic beta-blockers in the modern era: a meta-analysis of latest randomized controlled trials.** *Ann Noninvasive Electrocardiol.* 2013; 18:58-68.60.
24. Cook, R.C., Yamashita, M.H., Kearns, M. et al, **Prophylactic magnesium does not prevent atrial fibrillation after cardiac surgery: a meta-analysis.** *Ann Thorac Surg.* 2013; 95:533-541.



*“Comparison is the thief of joy.”*

**Unknown**

**AUTHORSHIP AND CONTRIBUTION DECLARATION**

Sr. #	Author-s Full Name	Contribution to the paper	Author=s Signature
1	Rehan Riaz	Mauscript Writing	<i>Rehan Riaz</i>
2	Nabeel Ahmad	Data collection	<i>Nabeel Ahmad</i>
3	Shahid Abbas	Statistical analysis	<i>Shahid Abbas</i>