

DR. WALI MUHAMMAD MAHERI

MBBS, M.Phil
Associate Professor Department of Physiology
Liaquat University of Medical
& Health Sciences Jamshoro

DR. MUHAMMAD ISHAQ

MBBS, FCPS(Pak)
FRCS(Ire) FRCS(Glasgow)
Department of Surgery
Jinnah Medical College
Peshawar.

DR. ISRAR AHMED AKHUND

MBBS, M.Phil
Professor of Physiology
Jinnah Medical College
Peshawar.

Dr. Muhammad Sabir

MBBS, M.Phil Professor of Anatomy
Jinnah Medical College
Peshawar.

ABSTRACT... Background: Despite advances in the prevention and treatment of hypertension over the past decade, hypertension remains an important public health challenge. Recent efforts to reduce the prevalence of hypertension have been focused on non-pharmacologic means, specifically diet. An increased intake of magnesium has been shown in some but not all studies to reduce blood pressure in patients with hypertension. Decreased Serum magnesium levels are associated with development of hypertension. **Aims:** This study was planned to investigate relation of serum magnesium with blood pressure in patients with mild uncomplicated hypertension. **Subjects & Methods:** Study was conducted at LUMHS hospital Jamshoro and DHQ hospital charsadda for the period of six months. Fifty known cases of uncomplicated mild hypertensive patients were selected, same number of healthy controls were also examined. **Results:** When results were summed up and test parameters were compared, it was seen that no significant differences were found in serum magnesium levels among both groups, when compared statistically. **Conclusion:** Finally we conclude that no relation of magnesium with hypertension was observed. **Suggestions:** Much more work on wide scale population may be needed to clarify the idea.

Key words: Magnesium, Hypertension.

INTRODUCTION

Ischaemic heart disease (IHD) is a leading cause of death in most of industrial and western world. A number of risk factors are associated with IHD, major being hypercholesterolaemia, hypertension, cigarette smoking, diabetes mellitus and stress and strain. Hypertension is an important accelerator of the atherosclerotic process and it frequently accompanies adult ischaemic heart disease¹. Hypertension is associated with an increased risk of clinical cardiovascular complication due to atherosclerosis and atheroma develops earlier in these patients. There are significant differences between the severity of the lesions of atherosclerosis in hypertensive and non-hypertensive subject². Magnesium is a biologically essential cation, which has recently received considerable attention in clinical medicine, especially with regard to the role of its depletion in cardiovascular pathophysiology³. Magnesium is the fourth most abundant cation in the body and the second most abundant intra cellular cation, next to potassium⁴. Some authors have shown recently an increasing interest in the effects of

calcium and magnesium on blood pressure⁵. Many reports have appeared in recent year discussing association between serum magnesium levels and hypertension. Accumulating evidence implicates a role of magnesium and pathophysiology of essential hypertension^{14,15} but its role in pathophysiology is still unclear¹⁶. Present study was carried out to investigate this burning issue. Study was aimed to investigate this relation.

SUBJECTS AND METHODS

Study was conducted in the department of cardiovascular diseases, LUMHS, Jamshoro and DHQ teaching Hospital, Charsadda, for the period of six months. Fifty known cases of mild uncomplicated hypertension were selected. Same numbers of healthy control were also selected. The information about name, age, sex, duration of their illness and blood pressure, smoking habits, and family history of cardiovascular disease were recorded. Patients with diuretic therapy, thyroid abnormalities, liver failure, renal failure or

Table. Comparison of mean values serum magnesium (mg/dl) in controls and hypertensive patients.

Group	Blood pressure (mmHg) Systolic/diastolic		Serum magnesium (mg/dl)	P-value
Controls (n=50)	126.00±_1.87	79.96±_1.39	2.94±_0.05	>0.5
Patients (n=50)	155.00±_1.52	103.62±_1.21	2.83±_0.10	>0.5

*Each value represents mean of individual observation
± indicates standard error of mean.*

alcoholics were excluded from the study. Fifty healthy controls were selected with no preexisting cardiovascular disease. A single casual supine blood pressure measurement was obtained by trained staff using a standard mercury sphygmomanometer according to W.H.O criteria. Blood samples of patients and healthy controls were drawn from antecubital vein by taking aseptic measures for determination of serum magnesium levels. Statistical analysis was done by student's t-test.

RESULT

When results were summed up and test parameters were compared it was seen that mean age of patients and controls was 40-50 years \pm 0.1 Serum magnesium levels (table) in control subjects were 2.94 ± 0.05 mg/dl, while mean serum magnesium levels in hypertensive subjects were 2.89 ± 0.03 mg/dl. The differences were found non-significant ($p > 0.5$) when evaluated statistically.

DISCUSSION

Investigations of the association between serum magnesium and blood pressure have yielded conflicting results. Hvarfner et al. found a positive association between serum magnesium and blood pressure in 58 hypertensive patients and 124 controls studied in Uppsala, Sweden. There was no difference between the relation identified in the hypertensive and control groups⁷. The relation between serum magnesium and blood pressure has also been examined by using data from national health and nutrition (NH&N) survey, no association was identified between serum magnesium and systolic blood pressure⁸. Similarly, data from a community based cross sectional study of elderly whites in Baltimore provided no evidence of an association between serum magnesium and blood pressure⁹. Peterson and co-workers reported a significant increase

in correlation between serum magnesium levels and systolic blood pressure¹⁰. Rinner et al studied Dutch population, and they found no relation between serum magnesium and blood pressure¹¹. Altura and Altura have reviewed the mechanism underlying in the relationship between magnesium and blood pressure. It has been postulated that if the concentration of extracellular magnesium is lowered, calcium influx is enhanced. There is relatively little information from both animal and human studies to indicate direct relationship between magnesium and blood pressure¹². Similarly, Herzog in 1995 also failed to establish any relationship between blood pressure and serum magnesium¹³.

In the present study, it was observed that the serum magnesium levels in patients with hypertension and in controls, are within normal limits. Therefore our study suggests no relation between serum magnesium levels and blood pressure in the patients of hypertension. Therefore, this study is in the favor of works done by Whelton et al, Rinner et al, Cappuccio et al, and Herzog et al^{6,9,11,13}. Our results are in contrast with the study done by Hvarfner et al. as they found high serum magnesium levels in hypertensive patients⁷. Also, study of Peterson and co-workers which reported decreased levels of serum magnesium in hypertensive patients, conflicts with our study¹⁰. Finally we suggest that further large scale studies on a large population are to be carried out to clarify idea.

Copyright© 4 Dec, 2010.

REFERENCES

1. Puska, P., Nissinen, A And Vartiainen, E. **Controlled. Randomized trial of the effect of dietary fat on blood pressure.** Lancet 2005;(I): 1-5.
2. Bondjers, G., Glukhova, M and Hansson, G.K.

- Hypertension and atherosclerosis.** *Circulation.* 2003;Pp.,12-16.
3. Ralston, M.A., murnane, M.R., Kelley, R.F., Altschuld, R.A., Unverferth, D.V. and leier, C.V. **Magnesium content of Serum circulation mononuclear cells, skeletal muscle, and Myocardium in congestive heart failure.** *Circulation.* 1989; 80:573-80.
 4. Rude, R.K. **physiology of magnesium metabolism and the important of magnesium in potassium deficiency.** *Am J Cardiol.*, 1989;63:31 G-34G.
 5. Kesteloo, H. and Joossents, J.V. **Relationship of dietary sodium, potassium, Calcium and magnesium with blood pressure.** *Hypertension.* 1988;12(6):564-9.
 6. Cappuccio, F.P., Markando, N.D., beynon, G.W., Shore, A.C. and McGregor **lack of oral magnesium and high blood pressure.** *BMJ.*, 1985;291:235-8.
 7. Hvarfner, A., Bergstrom, R. Morlin. C., Wide, L. and Ljunghall, R. **Relationship between magnesium metabolic indices and blood pressure in patients with essential hypertension, compared with a healthy Population.** *J. hypertens.*, 1987;5;451-6.
 8. Harlan, W.R., Hull, A.L., Schmouder, R.L., landis, Landis, J.R. Thompson, F.E. and larkin, F.A. **Blood pressure and nutrition in adults.** *Am. J.Epidimiol.*, 1984;120;17-28.
 9. Whelton, P.k. and klag, M.J. **Magnesium and blood pressure: review of the epidemiologic and clinical trial experience.** *Am.J. Cardiol.*, 1989; 63;26G-30G.
 10. Peterson, B., schroll, M., christiasen, C. and Transbol, I. **Serum and erythrocyte magnesium in normal elderly Danish people.** *Acta. Med. Scand.*, 1977;201:31-34.
 11. Rinner, M.D., Splient-Van-laar, L. and Kromhout, D. **Serum sodium, potassium, calcium and magnesium and blood pressure in a dutch population.** *J. hypertension.* 1989; 7 (12);977-81.
 12. Altura, B.M., Altura, B.T., Gebravold, A., Ising, H. and gunther, T. **Magnesium deficiency and hypertension co-relation between magnesium deficient diets and micro circulatory changes in situ.** *Science.*, 1984; 233:1315-7.
 13. Herzog, W.R (1995) **Reactivity of coronary arteries in relation to serum magnesium levels in patients with Hypertension.** *Circulation* 92:2622-26. Hatzistavri LS, Sarafidis PA(2009).
 14. **Oral magnesium supplementation reduces ambulatory blood pressure in patients with mild hypertension.** *Am. J. Hypertens.* 2002oct;(10):1070-5.
 15. Houston MC, Harper KJ: **Potassium, magnesium and calcium: their role in both the cause and treatment of Hypertension.** *J. Clin. Hypertens.* 2008 Jul; 10(7 Suppl 2):3-11.
 16. Paravicin TM, Yogi A, Mazur A: **Dysregulation of vascular TR PM7 and anmexin-1 is associated with endothelial in inherited hypomagnesemia.** *Hypertension:* 2009 Feb; 53(2): 432-9.

Article received on: 14/05/2010

Accepted for Publication: 04/12/2010

Received after proof reading: 00/00/0000

Correspondence Address:

Dr. Muhammad Ishaq
Jinnah Medical College
Peshawar.
info@jmcp.edu.pk

Article Citation:

Maheri WM, Ishaq M, Akhund IA, Sabir M. Serum magnesium and hypertension. *Professional Med J Mar* 2011;18(1):139-141.