

HNPCC

HEREDITARY NONPOLYPOSIS COLORECTAL CANCER IN PAKISTAN

ORIGINAL
PROF-1707

DR. MUTAHIR A. TUNIO, MBBS, FCPS

Radiation Oncology,
Sindh Institute of Urology & Transplantation (SIUT),
Karachi

DR. ALTAF HASHMI,

MBBS, MCPS, MS
Surgical Oncology,
Sindh Institute of Urology & Transplantation (SIUT),
Karachi

DR. REHAN MOHSIN, MBBS, FCPS

Surgical Oncology,
Sindh Institute of Urology & Transplantation (SIUT),
Karachi

Dr. Gohar Sultan, MBBS, FCPS

Surgical Oncology,
Sindh Institute of Urology & Transplantation (SIUT),
Karachi

ABSTRACT:... Hereditary nonpolyposis colorectal cancer (HNPCC) is the most common form of hereditary bowel cancer. Multiple generations are affected with colorectal cancer at relatively young age, between 25 and 45 years. **Objective:** To investigate the frequency of HNPCC in Pakistani population, due to the high incidence of colorectal cancer in younger Pakistani adults and prevalence of consanguinity in this region conducted this study. **Period:** November 2008 and January 2010. **Methodology:** Ninety histopathologically confirmed colorectal cancer patients between 12- 50 years and their families were interviewed using a detailed questionnaire. The questions about family history of colorectal cancer, history of other cancers, age at diagnosis and consanguinity were asked. The pedigrees were drawn for all families based on given information. Hospital records were also reviewed to confirm cancers reported in relatives. Amsterdam criteria was used to label a family as HNPCC. **Results:** Seventeen patients (18.9%) had one or more first or second degree relatives under age 50 years with colorectal cancers suggestive of HNPCC. Another 15 patients (16.7%) had first or second degree relatives with a family history of other extra-colonic cancers including ovarian, breast, endometrium, lung, parotid, brain and bladder cancer. Of these 30 patients (33.3%) reported that their parents were first degree cousins. **Conclusions:** 1. High frequency of HNPCC is seen in Pakistani population. 2. Higher proportion of colorectal cancer is seen in young Pakistanis. Strong prevalence of consanguineous marriages could be important factors for HNPCC occurrence in Pakistan. Future studies with large sample size along with genetic testing and screening programmes are warranted.

Key words: Hereditary non polyposis colorectal cancer, consanguinity, Pakistan

INTRODUCTION

Colorectal cancer (CRC) is amongst the common cancers worldwide. The incidence of colo-rectal carcinoma in Pakistan is similar to other Asian countries, but much lower than in the developed countries. The risk is equal in both sexes at present. However a 41 % rise in incidence was noted in the males between 1995 to 1999, which may indicate a higher risk in the males in future¹. Various factors and mechanisms have been discovered which play key role in the development of CRC^{2,3}. It has been thought that one-third of CRC patients have a genetic background. Hereditary nonpolyposis colorectal cancer (HNPCC) is a common autosomal dominant colorectal disorder with special clinicopathological features⁴. In clinic, it is diagnosed by Amsterdam criteria I:⁵ (1) three or more relatives with histologically confirmed colorectal cancer, one of whom is the first-degree to the other two; (2) colorectal cancer affecting at least two generations; and (3) one or more colorectal cancer cases diagnosed before the age of 50. Amsterdam criteria I is

too rigid, it excludes extra-colonic cancers and it may not be applicable for small families⁶. Alternatively, Amsterdam criteria II was defined:⁷ (1) at least three relatives must have a cancer associated with HNPCC (colorectal, endometrial, stomach, ovary, ureter or renal pelvis, brain, small bowel, hepatobiliary tract or skin tumors), (2) one must be first degree relative of other two, (3) at least two successive generations must be affected, (3) at least one of relative should have diagnosed before the age of 50, (5) familial adenomatous polyposis (FAP) should have been excluded in any relative with colorectal cancer and (6) tumor should be verified whenever possible.

The proportion of HNPCC varies by country, with rates ranging from 1 to 10% among all colorectal cancer patients (median 2-5%)⁸. Our previous studies mentioned that colorectal cancer patients under age 30 represent more than 30% of total colorectal cancer in our country^{9,10}.

Therefore, we conducted this pilot study to evaluate the frequency of HNPCC and associated colorectal and extra-colonic cancers according to Amsterdam criteria I and II in Pakistani patients and their families.

METHODOLOGY

Between November 2008 and January 2010, ninety histologically confirmed colorectal cancer patients aged between 16-50 years and their families were interviewed regarding risk factors for colorectal cancers after taking written consent. These patients represented about 60% of total colorectal cancer patients treated in same institute. The patients and families were interviewed by two of authors using a detailed questionnaire. The questions regarding family history of colorectal cancer, extra colonic cancer, age at time of diagnosis, symptoms, type of treatment and consanguinity (intercousin marriages) were asked. The pedigrees were drawn for the all families based on given information. All hospital histopathology reports were collected to confirm the diagnosis. Similarly to confirm the diagnosis in relative, the families were asked to provide the hospital record whenever possible. The Pakistani families have strong family ties and are aware of their relatives' cancer history.

The pedigrees were drawn using Smart Draw software. All data were analyzed by software SPSS version 16.0.

RESULTS

Ninety patients with confirmed colorectal cancer with median age 34.7 years (16-50) and their families were interviewed. Study population was predominantly males; 63 patients (70%). Rectosigmoid was found most frequent site (55.5%) of colorectal cancer in study population. 58 patients (64.4%) had no cancer family history and were considered sporadic. 32 patients (35.6%) had one or more first or second degree relative under age 45 years with colorectal cancers and extra-colonic cancers suggestive of HNPCC family (Table I).

The extra-colonic cancers were seen ovarian, breast, endometrium, lung, parotid, brain, bladder and esophageal cancer and most of these cancers occurred in paternal relatives (Table II).

Table-I. Characteristics of patients aged between 16-50 years with colorectal cancer

Variable	Patients (%)	
Median age (range)	34.7 years (16-50)	
Gender		
Male	63 (70%)	
Female	27 (30%)	
Site of primary tumor		
Proximal to splenic flexure	21 (23.4%)	
Descending colon	19 (21.1%)	
Rectosigmoid	50 (55.5%)	
Median number of first degree relatives (range)	10 (3-17)	
Relatives with cancer		
None	58 (64.4%)	
Relatives with colorectal cancers	17 (18.9%)	35.6%
Relatives with extra-colonic cancers	15 (16.7%)	
Consanguinity among first degree relatives (%)	30 (33.3%)	

Seventeen patients and their families fulfilled the Amsterdam criteria I for HNPCC for relatives in three generations were affected with colorectal cancers. Figure 1 shows the pedigree of a patient with a family history of colorectal cancers. The frequency of extra-colonic cancers were as; stomach cancer in 5 families (15.6%), non small cell lung cancers in 4 families (12.5%), breast cancer in 4 families (12.5%), endometrial cancers in 3 families (9.4%), brain astrocytomas in 3 families (9.4%), renal cell carcinoma in one family (3.1%), bladder cancer in one family (3.1%), ovarian cancer in one family (5.9%) and esophageal cancer in one family (5.9%). Figure 2 shows the pedigree of a patient with history of extra-colonic cancers.

Interestingly, 30 patients (33.3%) in study reported that their parents were first degree cousins. All patients with family history of consanguinity were relatively younger patients and had locally advanced cancers.

Further, ten families of colorectal cancers (11.1%) did not meet all of Amsterdam criteria, but they had colorectal cancers in two generations.

Table-II. Colorectal cancer patients with family history of one or more relative with extra-colonic cancer

Colorectal cancer patient (age and gender)	Relation with patient and age of diagnosis of cancer	Type of extra-colonic cancer
25 years, Female	Paternal grandfather (50 years)	Non-small cell lung cancer
30 years, Female	Sister (34 years) Brother (45 years) Mother (50 years)	Endometrial cancer Non-small cell lung cancer Breast cancer
30 years, Male	Sister (35 years) Mother (50 years)	Breast cancer Ovarian cancer
30 years, Male	Brother (32 years) Paternal uncle (50 years)	Stomach Non-small cell lung cancer
35 years, Male	Brother (45 years) Paternal uncle (41 years)	Stomach Stomach
25 years, Male	Paternal Grandmother (50 years) Maternal grandfather (52 years)	Parotid gland cancer Brain astrocytoma
45 years, Male	Maternal uncle (40 years) Maternal aunt (35 years)	Brain astrocytoma Breast cancer
30 years, Male	Paternal uncle (47 years) Maternal aunt (45 years) Maternal uncle (40 years) Maternal grandfather (50 years)	Lung cancer Renal cell cancer Non-small cell lung cancer Brain astrocytoma
32 years, Male	Father (50 years) Mother (47 years) Maternal uncle (41 years)	Non-small cell lung cancer Breast cancer Non-small cell lung cancer
40 years, Female	Sister (43 years)	Endometrial cancer
33 years, Male	Paternal uncle (60 years)	Bladder cancer
49 years, Female	Paternal grandmother (55 years)	Endometrial cancer
20 years, Male	Paternal uncle (49 years)	Stomach cancer
33 years, Female	Paternal grandfather (52 years)	Esophageal cancer
41 years, Male	Brother (44 years)	Stomach cancer

DISCUSSION

In 1913, Warthin described first time that some families with an excess of colorectal, endometrial and gastric

cancers. After 50 years, Lynch collected data that led to accurate description of these cancer-prone families¹¹. According to the absence or presence of extracolonic




Figure-2. The pedigree of 30 years old male patient with family history of predominant extra-colonic cancer (mother, paternal uncle, maternal uncle, paternal grandmother, and paternal grand father). (Note: box represents male and circle represents female)




Figure-1. the pedigree of a 32 years old male patient with a family history of colorectal cancers (maternal uncle, maternal grandfather). Other extra-colonic cancers were seen in his father, mother and maternal uncle. (Note: box represents male and circle represents female)

malignancies, these families were divided into Lynch syndrome I and Lynch syndrome II. Later, the Lynch syndrome was re-named as hereditary non-polyposis colorectal cancer (HNPCC)¹². HNPCC is an autosomal

dominant disease with special clinicopathological features; early onset (average <45ys), high frequency of cancers in right sided colon (60-70 %), excess of synchronous (tumors which appear on different sites of

colon within six months of surgery for first diagnosed colon cancer) (18.1 %) and metachronous (tumors which appear on different sites of colon after six months of surgery for first diagnosed colon cancer) (24.2 %) tumors and extra-colonic malignancies¹³. It is often diagnosed by Amsterdam criteria, but the criteria is too strict for families with small number of members but with strong genetic basis for colorectal cancer (CRC), in these cases Amsterdam criteria II is very useful¹⁴.

Presentation of Pakistani population at relatively younger age in our institute is alarming sign and was the basis of this study. Though this study was with small sample size, 35.6% of patients with typical HNPCC in these patients according to Amsterdam criteria explain higher proportion of HNPCC is our population. Another 11.1 % of families who did not meet all Amsterdam criteria may still have HNPCC and may need more investigations and screening.

CRC is multifactorial disorder and there can be several factors for its cause. One critic may rise about false positive rates in this study as affected relatives in families might have been exposed to environmental factors rather than genetic clustering of cancer. Explanation for this could be that family history of colorectal cancer is an important risk factor for CRC, and there is no evidence that familial clustering of CRC is affected by environmental factors¹⁵. There can be two other reasons for higher frequency of HNPCC is our population; (1) more prevalence of inter-cousin marriages (consanguinity) in Pakistan, as in our study 33.3% patients reported that their parents were first degree cousins and (2) Pakistani families are usually large, as our study showed median number of first degree relatives in a family were 10 (3-17). This factor may increase the relative risk of a positive HNPCC diagnosis, as suggested by Percesepe et al.¹⁶

Because early age is the feature of HNPCC, however cancer can also be detected in many families after age of 50 years, as we found in our study four relatives were aged above 50 years. This can also be explained by that, HNPCC is autosomal dominant disease and 50% of children with parents with HNPCC should be affected. However it takes years before the HNPCC is clinically manifested.¹⁷ One limitation of our study was that we did

not include the patients with colorectal cancers who were aged above 50 years.

Identification of HNPCC is not always simple because this disease can not be verified until many relatives have been affected with cancer and their exact age at time of diagnosis not exactly known to families (recall bias) are the key issues. Apart from Amsterdam criteria, genetic testing including microsatellite instability and mismatch repair gene mutations are required to confirm our results.^{18, 19} Already we are working on this and results will be provided in near future.

In conclusion, HNPCC is relatively more prevalent in Pakistan especially in colorectal cancer patients who present at early ages and this study highlights the importance for conducting future multi-institutional trial with large sample size incorporating genetic testing and screening of family members of young patients with colorectal cancer.

ACKNOWLEDGMENT

Authors are thankful to Professor Dr. Adibul Hasan Rizvi Director, Sindh Institute of Urology & Transplantation (SIUT), for his constant support as mentor and constant feedback.

Copyright© 20 June, 2011.

REFERENCES

1. Bhurgri Y, Bhurgri A, Nishter S, Ahmed A, Usman A, Pervez S, et al. Pakistan-country profile of cancer and cancer control 1995-2004. *J Pak Med Assoc.* 2006;56:124-30.
2. Markowitz S. **DNA repair defects inactivate tumor suppressor genes and induce hereditary and sporadic colon cancers.** *J Clin Oncol* 2000, 18:75-80.
3. Boland CR, Goel A. **Microsatellite instability in colorectal cancer.** *Gastroenterology* 2010;138:2073-87.
4. Jasperson KW, Tuohy TM, Neklason DW, Burt RW. **Hereditary and familial colon cancer.** *Gastroenterology* 2010; 138:2044-58.
5. Vasen HF, Mecklin JP, Khan PM, Lynch HT. **The International Collaborative Group on Hereditary Non-Polyposis Colorectal Cancer (ICG-HNPCC).** *Dis Colon Rectum.* 1991;34:424-5.

6. Walsh MD, Buchanan DD, Cummings MC, Pearson SA, Arnold ST, Clendenning M, et al. **Lynch syndrome-associated breast cancers: clinicopathologic characteristics of a case series from the colon cancer family registry.** Clin Cancer Res 2010;6:2214-24.
7. Vasen HF, Watson P, Mecklin JP, Lynch HT. **New clinical criteria for hereditary nonpolyposis colorectal cancer (HNPCC, Lynch syndrome) proposed by the International Collaborative group on HNPCC.** Gastroenterology. 1999;116:1453-6.
8. Soliman AS, Bondy ML, Levin B, Khaled H, Hablas A, Ismail S, et al. **Familial aggregation of colorectal cancer in Egypt.** Int J Cancer 1998;77:811-6.
9. Abbas A, Tunio MA, Ali N. **Neoadjuvant chemo-irradiation in un-resectable carcinoma of rectum.** J Ayub Med Coll Abbottabad. 2004;16:24-9.
10. Tunio M, Rafi M, Maqbool A, Haque A. **Virtual simulation and treatment verification-merits and demerits: Experience at Sindh Institute of Urology and Transplantation (SIUT).** Journal of Radiotherapy in Practice 2009; 8:131-36.
11. Marra G, Boland CR. **Hereditary nonpolyposis colorectal cancer: the syndrome, the genes, and historical perspectives.** J Natl Cancer Inst 1995;87:1114-25.
12. Nagasaka T, Rhee J, Kloor M, Gebert J, Naomoto Y, Boland CR, et al. **Somatic hypermethylation of MSH2 is a frequent event in Lynch Syndrome colorectal cancers.** Cancer Res 2010;70:3098-108.
13. Kohlmann W, Gruber SB. **Hereditary Non-Polyposis Colon Cancer.** In: Pagon RA, Bird TC, Dolan CR, Stephens K, editors. GeneReviews. Seattle (WA): University of Washington, Seattle;1993-2004.
14. Mvundura M, Grosse SD, Hampel H, Palomaki GE. **The cost-effectiveness of genetic testing strategies for Lynch syndrome among newly diagnosed patients with colorectal cancer.** Genet Med 2010;12:93-104.
15. Kotake K, Koyama Y, Nasu J, Fukutomi T, Yamaguchi N. **Relation of family history of cancer and environmental factors to the risk of colorectal cancer: a case-control study.** Jpn J Clin Oncol 1995;25:195-202.
16. Percesepe A, Borghi F, Menigatti M, Losi L, Foroni M, Di Gregorio C, et al. **Molecular screening for hereditary nonpolyposis colorectal cancer: a prospective, population-based study.** J Clin Oncol 2001;19:3944-50.
17. Benatti P, Roncucci L, Ganazzi D, Percesepe A, Di Gregorio C, Pedroni M, et al. **Clinical and biologic heterogeneity of hereditary nonpolyposis colorectal cancer.** Int J Cancer 2001;95:323-8.
18. Pedroni M, Sala E, Scarselli A, Borghi F, Menigatti M, Benatti P, et al. **Microsatellite instability and mismatch-repair protein expression in hereditary and sporadic colorectal carcinogenesis.** Cancer Res 2001;61:896-9.
19. Russo A, Sala P, Alberici P, Gazzoli I, Radice P, Montefusco C, et al. **Prognostic relevance of MLH1 and MSH2 mutations in hereditary non-polyposis colorectal cancer patients.** Tumori 2009;95:731-8.

Article received on: 11/10/2010

Accepted for Publication: 20/06/2011

Received after proof reading: 12/08/2011

Correspondence Address:

Dr. Mutahir A. Tunio, MBBS, FCPS
 Radiation Oncology
 Sindh Institute of Urology & Transplantation (SIUT),
 Karachi
 drmutahirtonio@hotmail.com

Article Citation:

Tunio MA, Hashmi A, Mohsin R, Sultan G. Hereditary nonpolyposis colorectal cancer in Pakistan: results of a pilot study. Professional Med J Sep 2011;18(3): 344-349.

PREVIOUS RELATED STUDIES

- Muhammad Akram, Wasim Amer, Javaid Iqbal. Colorectal cancer; overall survival and its prognostic factors, a retrospective analysis from 10 year data form JHL. Professional Med J Dec 2009; 16(4):492-498.